

INDEX

- Abortion Act in Britain, 103, 281
 abortion trends, legal, 281
 accident and emergency department, 240
 care in, 44
 general failings in, 46
 recommendations, 45
 acquired cardiac disease, causes of
 aneurysm, 140
 cardiomyopathy, 140
 myocardial infarction, 140
 acquired heart disease cases, deaths from
 aortic aneurysm and dissection, 145
 cardiomyopathy, 146
 coronary artery dissection, 144
 hypertensive heart disease, 147
 ischaemic heart disease, 144
 myocardial fibrosis, 147
 acute fatty liver, maternal deaths from, 120
 acute respiratory distress syndrome (ARDS), 80
 Advanced Life Support in Obstetrics, 100. *See also*
 amniotic fluid embolism
 age
 maternities by, 284, 285
 risk factor for maternal deaths, 51
 alcohol related deaths, 162
 midwives key recommendations for, 259, 260
 See also drug/alcohol related deaths
 amniotic fluid embolism
 cases
 clinical features of, 99
 speed of collapse, 98
 diagnosis (case history), 228
 diagnosis criteria
 acute hypotension/cardiac arrest, 100
 acute hypoxia, 100
 coagulopathy, 100
 findings for 2000–02, 97
 recommendations
 individual practitioners, 96
 pathology, 96
 service provision, 96
 substandard care for, 100
 See also thromboembolism
 amniotic fluid register, 100
 anaesthesia, recommendations for
 individual practitioners, 122
 service provision, 122
 anaesthetic maternal deaths
 causes of
 anaphylaxis, 128
 aspiration of gastric contents, 128
 isolated sites, 127
 lack of appreciation of severity of illness, 129
 lack of multidisciplinary cooperation, 129
 management of haemorrhage, 131
 misplaced tracheal tubes, 126
 perioperative care, 130
 history of, 123
 rapid-sequence induction technique, 124
 regional anaesthesia, use of, 124
 tracheal intubation for caesarean section, 124
 numbers of, 124
 anaphylaxis, 128
 aneurysms, autopsy report of, 229
 antenatal care
 improvements in, 57
 midwifery practice and, 245
 poor attendance at antenatal clinic, 40
 See also substandard care
 antenatal prophylactic and therapeutic doses,
 guidelines for, 69
 antepartum deaths, 67
 antihypertensive treatment, 83
 aortic aneurysm
 autopsy report of, 229
 during pregnancy, 145
 aortic dissection, diagnosis of, 145
 asylum seekers, maternal deaths among, 47
 autopsy on maternal death, assessment criteria for,
 224
 autopsy report
 added value of, 227
 amniotic fluid embolism, 228
 aneurysms, 229
 cardiac disease, 228
 congenital heart disease, 230
 diabetic pregnant mothers, 232
 dissecting coronary artery aneurysms, 230
 epilepsy, 231
 Ogilvie's syndrome, 232
 suicide, 231
 findings for 1973–75, 221
 findings for 1997–99, 222
 findings for 2000–02, 223
 in triennial reviews of enquiry, 222
 on maternal deaths, 221
 clinical correlation, 226
 history of, 221
 substandard autopsies, 225
 performance of (case history), 225
 quality of, 224
 bacterial endocarditis causing maternal deaths
 (indirect), 148
 bacterial infection
 meningococcus meningitis, 188
 pneumococcus meningitis, 188
 staphylococcal septicaemia, 188

- beta-hCG testing for early pregnancy deaths, 106
- birth rates and fertility trends, 283
- blood diseases, 194
 - causes of
 - failure in severity assessment, 192
 - failure to identify abnormal test results, 193
 - poor communication with pregnant women, 192
 - haemolytic uraemic syndrome, 192
 - thrombotic thrombocytopenic purpura (TTP), 192
- blood loss in pregnant woman. *See* haemorrhage
- blood pressure control, 83, 84, 130
- blood transfusion refusal leading to death, 91.
 - See also* postpartum haemorrhage
- body mass index (BMI), 50, 291. *See also* obesity
- booking visit by maternity staff, 20, 307
 - booking referral letter, 21
 - risk assessment framework, 21
- bowel perforation, 232
 - maternal deaths from, 120
- breast examination, 201
- caesarean section
 - 2004 guideline on, 38
 - by place of delivery
 - perimortem, 39
 - postmortem, 39
 - causing genital tract sepsis, 114
 - in 2000-02 (Hospital Episode Statistics (HES) data), 38
 - risk factors for deaths after, 69
 - age factor, 70
 - cerebral vein thrombosis, 70
 - diagnosis, failure of, 70
 - family history, 70
 - late deaths, 70
 - obesity factor, 70
 - thromboprophylaxis, adequacy of, 70
 - types, RCOG definition of
 - elective, 38
 - emergency, 38
 - perimortem, 38
 - postmortem, 38
 - scheduled, 38
 - urgent, 38
- cancer deaths (maternal)
 - causes/types of deaths, 199
 - lack of investigation in pregnancy
 - abdominal pain and vomiting, 201
 - chronic chest/abdominal/back pain, 201
 - epigastric pain, 201
 - multidisciplinary care, 202
 - poor communications between professionals, 200
 - prepregnancy counselling, 200
 - recommendations for
 - individual practitioners, 197
 - service provision, 197
 - social labelling affecting diagnosis, 202
- cancer diagnosis in pregnancy, delay in
 - lack of investigation, 201
 - poor communications between professionals, 200
 - social labelling, 202
- cancer in pregnancy, study on, 198
- cardiac disease
 - accuse of maternal death, 228
 - post-delivery care for cardiac patients, 149
 - See also* heart diseases
- cardiac disease causing maternal deaths
 - acquired causes, 140
 - congenital causes, 140
 - findings for 2000–02, 140
 - history of (1952–2002), 138
 - recommendations for
 - guidelines and protocols, 137
 - individual practitioners, 137
 - service provision, 137
- cardiomyopathy
 - autopsy report of, 229
 - causing maternal mortality
 - dilated, 146
 - hypertrophic obstructive, 146
 - peripartum, 146
 - diagnosed at autopsy, 231
 - diagnosis of, 146
- CEMACH for England and Wales, 315–318
 - maternal death enquiry, 3
 - use of vignettes, 4
 - See also* confidential enquiries into maternal deaths, 1952 onwards trends and development
- central nervous system disease, 188, 194
 - cerebral thromboembolism, 187
 - intracerebral haemorrhage, 187
 - intracranial haemorrhage, 186
 - subarachnoid haemorrhage, 186
- cerebral thromboembolism (case history), 187
- cervical intraepithelial neoplasia (CIN), case history of, 200
- child protection issues, 170, 180
- circulatory system, diseases of
 - ruptured iliac artery, 193
 - ruptured pelvic vein, 193
 - ruptured spleen, 193
- clinical care for pregnant women, 19
- clinical correlation, 226
- clinical standards setting for maternity services
 - Clinical Negligence and Other Risks Indemnity (CNORIS), 6
 - Clinical Negligence Scheme for Trusts (CNST), 5
 - Clinical Resource and Audit group (CRAG), 5

- Clinical Standards Board for Scotland (CSBS), 6
- Scottish Intercollegiate Guidelines Network (SIGN), 5
- clinicopathological correlation, 223
- coincidental deaths, 157, 160, 177
 - causes for late deaths, 217
 - during pregnancy
 - domestic violence and murder, 210
 - road traffic accidents, 208, 209
- coincidental deaths from domestic violence
 - findings for 2000–02
 - age at death, 210
 - characteristics of women, 212
 - murder by family members, 211
 - number of deaths, 209
 - number of deaths, 211
 - secret keeping, 210
 - social structure of victims, 210
 - recommendations for
 - individual practitioners, 207
 - service provision, 207
 - training, 208
- confidential enquiries into maternal deaths, 1952
 - onwards trends and development, 296, 299
- avoidable factors and substandard care, 306
- booking arrangements, 307
- changes in classification, 300
- completeness of ascertainment, 301
- international comparisons, 310
- maternal morbidity, 310
- report scope and agenda, 298
- social factors, 309
- See also* CEMACH for England and Wales
- Confidential Enquiry into Maternal and Child Health. *See* CEMACH for England and Wales
- congenital heart disease
 - autopsy report, 230
 - deaths from, 140
 - delay in obtaining termination of pregnancy, 143
 - from postpartum haemorrhage (PPH), 141
 - with Eisenmenger's syndrome, 143
 - with ventricular septal defect (VSD), 141
 - pulmonary hypertension cause of, 140
 - See also* heart diseases
- cystic medial necrosis, 230
- deep venous thrombosis (DVT), 63
- delivery, care for
 - antenatal care, 35
 - delivery, types of, 37
 - place of delivery, 37
- delivery types
 - caesarean section, 38
 - vaginal delivery, 38
- deprivation classification, 48
- diabetes mellitus, autopsy report of, 232
- diagnosis failure for caesarean section, 70
- direct causes for late deaths, 216
- direct deaths (maternal), 9
 - acute fatty liver, 120
 - amniotic fluid embolism, 96
 - anaesthesia and. *See* anaesthetic maternal deaths
 - bowel perforation, 118, 120
 - early pregnancy deaths, 102, 105
 - genital tract sepsis, 109, 112
 - genital tract trauma, 118, 120
 - obstetric haemorrhage, 86
 - pre-eclampsia and eclampsia, 79
 - thromboembolism, 61, 243
 - thrombosis, 61
 - See also* indirect deaths (maternal)
- dissecting coronary artery aneurysms, maternal deaths from, 230
- disseminated intravascular coagulation (DIC), 81, 109, 226
- domestic violence
 - coincidental deaths and. *See* coincidental deaths from domestic violence
 - indicators related to maternity care, 213
 - key recommendations for midwives, 258, 259
 - maternal deaths from, 49, 207
- drug/alcohol related deaths
 - education/training
 - maternity care staff, 174
 - primary care staff, 174
 - findings for 2000–02
 - bereavement or loss, 179
 - causes of death, 178
 - child protection issues, 180
 - coexisting mental health problems, 179
 - planning of pregnancies, 181
 - polydrug users, 180
 - reproductive choices, 181
 - services required, 179
 - substance misuse, 178, 180
 - timing of death, 177
 - individual practitioners, 174
 - midwives key recommendations for, 259, 260
 - problem drug use
 - multi-agency approach, 176
 - prevalence and patterns of, 175
 - socio-economic deprivation factor, 176
 - service delivery, 174
- drug/alcohol use in pregnancy, 182, 259
- early pregnancy deaths, 105
 - beta-hCG testing for, 106
 - recommendations for
 - individual practitioners, 102
 - service provision, 102
 - training, 102

- ectopic pregnancy, maternal deaths from, 105.
 - See also* early pregnancy deaths
- endocrine, metabolic and immunity disorders, 194
 - diabetes, 190
 - phaeochromocytoma, 191
- England and Wales 1997–2002 report on pregnant women mortality
 - comparison with non pregnant women mortality, 272
 - computational methods, 273
 - results, 274
- epidemiology trends, reproductive
 - birth rates and fertility trends, 283
 - delivery complications, 290
 - delivery mode, 289
 - legal abortion trends, 281
 - maternal health aspects
 - alcohol, 292
 - blood pressure, 291
 - contraceptive pill usage, 292
 - ethnic populations health, 294
 - obesity, 291
 - postnatal depression and other psychological illness, 293
 - smoking, 292
 - maternities and estimated pregnancies, 279
 - maternities by
 - age, 284, 285
 - ethnic origin, 286, 287
 - marital status, 286
 - multiplicity, 288
 - National Statistics socio-economic group, 287, 288
 - parity, 286
- epilepsy
 - autopsy reports of, 231
 - causing maternal deaths, 187
- ethnic origin
 - maternities by, 286, 287
 - risk factor for maternal deaths, 45, 244
- fertility trends, 283
- fluid management in pregnant women, 84.
 - See also* blood pressure control
- fungal infection from *aspergillosi*, 189
- gastrointestinal system, diseases of
 - intestinal obstruction, 191
 - pancreatitis, 191
 - peritonitis, 191
 - ruptured oesophagus, 191
- general practitioners role in maternal death prevention, 58
- genital tract sepsis
 - after surgery, 114
 - before delivery, 112
 - causing maternal mortality, history of, 109
 - diagnosis for, 114
 - findings for 2000–02, 111
 - intrapartum deaths, 113
 - sepsis after vaginal delivery, 113
 - sepsis before delivery, 112
 - investigating, 115
 - maternal mortality rates (1847–2002), 110
 - recommendations for
 - education and training, 109
 - individual practitioners, 109
 - service provision, 109
 - signs of critical illness
 - biochemical, 116
 - physiological, 116
 - symptoms of
 - abdominal pain, 116
 - diarrhoea, 116
 - vomiting, 116
 - treatment of suspected, 115
- genital tract trauma deaths
 - history of, 118
 - maternal deaths from, 120
- haemorrhage
 - intracerebral, 187
 - intracranial, 186
 - risk factors for, 87
 - subarachnoid, 186
- haemorrhage (from anaesthesia), management of
 - blood transfusion, 131
 - cell saver, 132
 - consultant obstetric anaesthetists, availability of, 131
 - direct arterial pressure monitoring, 131
 - postoperative care, 132
- haemorrhage management
 - conventional, 239
 - with intensive care, 238, 239
- haemorrhage problems
 - placenta praevia, 90
 - postpartum haemorrhage, 91
 - resuscitation, quality for, 89
 - substandard care for, 89, 92
- health professionals
 - responsibilities, 22
 - training, 23
- healthcare workers role on maternal deaths report, 53
- heart disease
 - acquired heart disease cases, 144–147
 - congenital, 140, 141, 143, 230
 - hypertensive, 147
 - ischaemic, 144
- histology
 - failure of, 227
 - need for, 225

- human chorionic gonadotrophin (hCG), 102
- hypertensive heart disease, medical complications in, 147
- ICD, 300
- ICD10, 272
- immigrants (recent), maternal deaths among, 47
- immunohistochemistry, use of, 228
- in vitro fertilisation (IVF) for infertility. *See* infertility treatment causing maternal deaths
- indirect causes for late deaths, 216
- indirect deaths (maternal), 9
 - aspects of, 195
 - causes, 186, 244
 - central nervous system diseases, 186
 - chromosomal abnormalities, 195
 - circulatory system diseases, 193
 - diseases of blood, 192
 - endocrine, metabolic and immunity disorders, 190
 - epilepsy, 187
 - from cardiac disease, 137
 - from psychiatric illness, 244
 - gastrointestinal system, 191
 - infectious diseases, 188
 - recommendations for
 - individual practitioners, 183
 - service provision, 183
 - renal system diseases, 193
 - respiratory system diseases, 190
 - sudden adult death syndrome, 147
 - See also* direct deaths (maternal)
- infanticide
 - and filicide, 165
 - associated with maternal mental illness, 170
- infectious diseases
 - bacterial infection, 188
 - fungal infection, 189
 - viral infection, 189
- infertility treatment causing maternal deaths, 51
- intensive care for pregnant woman
 - accident and emergency departments, 240
 - early referral, 237, 238
 - findings for 2000–02, 236
 - history of
 - acute respiratory distress syndrome in adults (ARDS), 236
 - intensive care medicine, 234
 - intensive care units, 235
 - microprocessor-controlled medical devices, 235
 - in obstetric theatres, 238
 - pulmonary embolus, 240
 - recommendations for
 - individual practice, 234
 - service provision, 234
 - severity of illness, assessing
 - massive haemorrhage management, 238, 239
 - metabolic acidosis, 237
 - tachypnoeic, 237
- intensive care, features of, 241
- International Classification of Diseases, 300
- interpreting services, midwifery and, 251
- intrapartum deaths (streptococcal infection), 113
- ischaemic heart disease causing maternal mortality, 144. *See also* heart disease
- late deaths, 177
 - indirect deaths, 157
 - interval between delivery and maternal death, 216
 - maternal death, 10
 - See also* early pregnancy deaths
- late deaths, findings for 2000–02, 215
 - direct causes, 216
 - indirect causes (ONS record linkage study), 217
 - late coincidental deaths
 - causes, 217
 - ONS record linkage study, 218
- local maternity services, recommendations for, 18
- magnesium sulphate, 79. *See also* pre-eclampsia and eclampsia
- malignancy causing maternal deaths. *See* cancer deaths (maternal)
- Managing Obstetric Emergencies and Trauma (MOET), 100. *See also* amniotic fluid embolism
- marital status
 - maternities by, 286
 - risk factor for maternal deaths, 52
- maternal death enquiry (2000–02), key findings for, 30
- maternal deaths risk factors
 - domestic violence, 26
 - inter-agency communications, lack of, 27
 - inter-professional communications, lack of, 27
 - late booking or poor attendance, 26
 - minority ethnic groups, 26
 - obesity, 26
 - poor communities, 26
 - social disadvantage, 26
 - suboptimal clinical care, 26
 - substance abuse, 26
- mortality rates, 26
- See also* maternal deaths; maternal deaths enquiry
- maternal death problems
 - multidisciplinary care for, 42
 - recommendations for
 - birth, emergency services for, 22
 - booking visit, 20

- maternal death problems (*Cont.*)
 - clinical care, 19
 - complicated pregnancies, care for, 21
 - health professionals, 22
 - local maternity services, planning, 18
 - maternal deaths, reporting, 24
- maternal death report, key recommendations for
 - individual practitioners, 58
 - research, 57
 - service provision, 57
- maternal deaths, 30
 - analysis using 2004 English Indices of Deprivation. *See* Deprivation classification
 - autopsy, 224
 - by classification of woman
 - husband's occupation-based, 48
 - single mothers, 48
 - by gestation, type of death, 35
 - by mode and place of delivery, 37
 - by type of antenatal care, 36
 - direct. *See under* direct deaths (maternal)
 - estimated rates by type and ethnic group, 46
 - from cardiac disease, 138
 - acquired, 140
 - causes of, 141
 - congenital, 140
 - from drug and alcohol, 174
 - from eclampsia and pre-eclampsia (1988–2002), 81
 - from ectopic pregnancies, 105
 - from haemorrhage (1952–2002), 86
 - from haemorrhage, fall in
 - better surveillance of ill women, 86
 - coagulopathy, management of, 87
 - flying squad, use of, 87
 - hospitalisation for delivery, 86
 - from malignancy (cancer). *See* cancer deaths (maternal)
 - from miscarriage, 103
 - from psychiatric causes. *See* psychiatric cause for maternal deaths
 - from terminations of pregnancy, 103
 - histology, 227
 - in UK (1985–2002), number of
 - maternal death enquiry, reported to, 28
 - Registrars General, reported to, 28
 - indirect. *See under* indirect deaths (maternal)
 - pathology recommendations for, 221
 - rate calculation, 45
 - reporting for, 24
 - risk factors
 - access to care, 40
 - age, 51
 - asylum seekers and recent immigrants, 47
 - care in accident and emergency departments, 44
 - deprivation classification, 48
 - domestic violence, 49, 244
 - ethnicity, 45
 - infertility, 51
 - lack of interprofessional communications, 245
 - late booking or poor attendance, 244
 - marital status, 52
 - minority ethnic groups, 244
 - multiple pregnancies, 51
 - National Statistics Socio-Economic classification, 48
 - obesity, 50, 244
 - parity, 51
 - poor communities, 244
 - social disadvantage, 244
 - suboptimal clinical care, 245
 - substance abuse, 244
 - substance misuse, 50
 - substandard care, 42
 - translation, 47
 - with substandard care, 43
- maternal deaths before delivery
 - cardiac disease, 35
 - ectopic pregnancy, 35
 - pulmonary thromboembolism, 35
- maternal deaths, causes of, 14, 16, 26, 227
 - direct causes
 - eclampsia, 15
 - postpartum haemorrhage, 15
 - pre-eclampsia, 15
 - puerperal sepsis, 15
 - indirect causes
 - infectious diseases, 15
 - vector born diseases, 15
 - See also* maternal deaths, specific causes of
- maternal deaths enquiry
 - aims and objectives, 4
 - background, 3
 - benefits of, 17
 - clinical governance, contribution to, 5
 - clinical standards setting, role in, 5
 - confidential enquiries (1952 onwards trends and development), 296, 299
 - avoidable factors and substandard care, 306
 - booking arrangements, 307
 - changes in classification, 300
 - completeness of ascertainment, 301
 - international comparisons, 310
 - maternal morbidity, 310
 - report scope and agenda, 298
 - social factors, 309
 - role of CEMACH Regional Managers, 12
 - role of Office for National Statistics (ONS) in, 12
 - study methodology, 6
 - use by healthcare trusts
 - at local level, 5
 - at national level, 5
- maternal deaths, specific causes of, 31

- direct deaths
 - amniotic fluid embolism, 32
 - anaesthetic, 32
 - ectopic, 32
 - haemorrhage, 32
 - hypertension, 32
 - sepsis, 32
 - thromboembolism, 32
- indirect deaths
 - cardiac, 32
 - CNS haemorrhage, 32
 - epilepsy, 32
 - GI, 32
 - infections, 32
 - psychiatric, 32
 - respiratory, 32

See also maternal deaths, causes of
- maternal disabilities, causes of, 14
- maternal morbidity, 267, 270
 - severe, 310
- maternal mortality
 - by maternal age, 51
 - definitions
 - coincidental (fortuitous), 10
 - direct maternal deaths, 9
 - indirect maternal deaths, 9
 - late, 10
 - pregnancy-related death, 10
 - definitions, used in maternal death enquiry (UK), 11
 - from amniotic fluid embolism (1961–84), 97
- maternal mortality rate
 - calculation
 - deaths from obstetric causes, 13
 - estimated pregnancies, 13
 - number of maternities, 13
 - calculation (UK)
 - deaths reported to Enquiry, 27
 - official death certification, 27
 - estimation of, 10
 - for multiple births (1997–2002), 51
 - from genital tract sepsis, 110
 - in UK (1985–2002), 28
 - in UK (2000–02), 29
- maternal mortality ratio (MMR), 7, 10
- maternities by
 - age, 284, 285
 - ethnic origin, 286, 287
 - marital status, 286
 - multiplicity, 288
 - National Statistics socio-economic group, 287, 288
 - parity, 286
- mental health, 255
 - problems associated with substance misuse, 179
 - recommendations for midwives, 257
- midwifery practice, findings and recommendations
 - in, 243
 - antenatal care, 245
 - as an individual practitioner, 248
 - at booking, 246
 - domestic violence recommendations, 259
 - for drug and/or alcohol related deaths, 260
 - minority ethnic groups, 250
- midwifery practice issues, from 2000–02 findings, 249
 - advocacy, 262
 - challenges for future care delivery and improvement, 262
 - continuing professional development, 261
 - developing care to meet individual needs
 - advocacy, 253
 - deprivation and social exclusion, 250
 - domestic violence, 258
 - drug and/or alcohol misuse, 259
 - following up women who failed to attend antenatal clinics, 253
 - interpreting services, 251
 - mental health, 255, 257
 - minority ethnic groups, 250, 251
 - normal delivery practices, recognising deviations from, 261
 - professional accountability
 - communication, 260
- miscarriage causing maternal deaths, 106
- misplaced tracheal tubes causing anaesthetic deaths, 126
- morbidity. *See* maternal morbidity
- mortality in pregnant and non pregnant women, in England and Wales 1997–2002, 272
- mortality rates
 - by cause of maternal deaths, 31
 - calculation by method of delivery, 38
 - from haemorrhage by cause
 - amniotic fluid embolism, 88
 - eclampsia, 88
 - placenta increta at termination of pregnancy, 88
 - ruptured uterus, 88

See also maternal mortality
- multi-agency management (for drug/alcohol users), 176
- multidisciplinary care for cancer in pregnancy by
 - anaesthetists, 202
 - midwives, 202
 - nurses, 202
 - obstetricians, 202
 - oncologists, 202
 - paediatricians, 202
 - radiologists, 202
 - woman's dignity protection, 203

- multiple pregnancies
 - maternities by, 288
 - risk factors for maternal deaths, 51
- murder of pregnant woman
 - by family members, 211
 - circumstances of, 213
- myocardial fibrosis, diagnosis of, 147
- myocardial infarction. *See under* acquired heart disease cases, causes of

- National Amniotic Fluid Embolism Register (UK).
 - See* amniotic fluid register
- National Institute for Clinical Excellence (NICE)
 - recommendations, 239
- National Statistics Socio-Economic Classification (NS SEC), 48
- near misses, 267

- obesity, 291
 - causing aspiration of gastric contents, 128
 - for death after caesarean section, 70
 - maternal deaths cause
 - body mass index (BMI) of 30, 50
 - body mass index (BMI) of 35, 50
 - midwifery recommendations, 255
 - risk factor for thromboembolism, 66, 68
- obstetric (complicated pregnancies), 21
- obstetric anaesthetists, responsibilities of, 129
- obstetric haemorrhage
 - management of, 92
 - recommendations for
 - individual practitioners, 86
 - service provision, 86
- obstetric theatres, 238
- Ogilvie's syndrome, pathology report of, 232
- ONS (Office for National Statistics)
 - for maternal death enquiry, 12
 - study on maternal death linkage, 31
 - deaths before delivery, 35
 - suicide (psychiatric cause), 33
- oral contraception causing maternal deaths, 67

- parity
 - maternities by, 288
 - risk factor for maternal deaths, 51
- pathology recommendations for pregnant woman, 221
- perinatal mental illness leading to suicide, 167
- perinatal psychiatric disorder, 153
- perinatal psychiatric services, specialist, 154
- perioperative care (anaesthesia services), 130
- phaeochromocytoma, 191
- placenta praevia diagnosis, 90
- placental abruption, features of, 90
- polydrug users, 180
- postmortem findings correlation with clinical events, 226
- postnatal mental illness
 - biological factors (genetic and neuroendocrine), 153
 - psychosocial factors, 153
- postnatal psychiatric disorder, 154
- postpartum deaths, 159
- postpartum haemorrhage (PPH), 87, 91, 227
- pre-eclampsia and eclampsia
 - 1952–96 deaths from, 80
 - consequences of, 84
 - deaths from
 - anti-hypertensive therapy, failure of, 81
 - clinical care, lack of, 81
 - disseminated intravascular coagulation (DIC), 81
 - from ARDS, 81
 - intracranial haemorrhage, 81
 - multi-organ failure, 81
 - systolic blood pressure, 82
 - treatment
 - anticonvulsant drug for, 79
 - guidelines for, 79
 - intensive care for, 79
- pregnancy deaths in early stages. *See* early pregnancy deaths
- pregnancy-related death, 10
- pregnant woman
 - characteristics of
 - late bookers, 40, 41
 - non-attenders at clinic, 40, 41
 - poor attenders at clinic, 40
 - unknown to antenatal maternity services, 41
 - concealing pregnancies, 42
 - mortality, in England and Wales 1997–2002
 - comparison with non pregnant women mortality, 272
 - computational methods, 273
 - results, 274
 - poor care for (examples), 45
- prepregnancy counselling about cancer, 200
- psychiatric care during index maternity, 163
- psychiatric cause for maternal deaths, 113
 - age at death, 161
 - child protection issues, 170
 - clinical isolation and communication, 170
 - communication between professionals and services, 165
 - contact with psychiatric services, 163
 - diagnosis of, 162
 - ethnicity, 161
 - findings for 1997–99, 166
 - findings for 2000–02, 156
 - illicit drug overdoses, 157
 - infanticide and filicide, 165
 - living children, number of, 165
 - maternal suicide, 154
 - perinatal psychiatric disorder, 153

- physical illness (intravenous drug user), deaths from, 158
- physical illness associated with psychiatric disorders, 159
- record-linkage study by ONS, 155
- report for 1997–99, 155
- risk of recurrence, 154
- social characteristics, 161
- specialist perinatal psychiatric services, 154
- substance misuse, 162
- suicide, 152
- timing of deaths, 158
 - from drug overdose, 160
 - from suicide, 160
- psychiatric deaths, 171
- psychiatric disorder
 - case history of, 168
 - clinical symptoms of, 169
 - diagnosis of, 162
 - misattribution of symptoms to, 169
 - perinatal, 153
 - postnatal, 154
 - risk detection, 167
 - risk identification, 164
 - risk management, 164, 167
- psychiatric illness, as indirect deaths cause, 244
- psychiatric management, 154
- psychiatric services
 - and professionals, lack of communication, 165
 - contact with, 164
 - history of, 163, 164
- puerperal sepsis. *See* genital tract sepsis
- pulmonary embolism
 - diagnosis for, 68
 - number of deaths from, 65
 - See also* amniotic fluid embolism; thromboembolism
- pulmonary embolus
 - autopsy reports of, 226
 - diagnosis of, 240
- pulmonary hypertension, 140
 - hypertensive changes in pregnancy (case history), 230
- renal disease, 195
 - chronic renal failure, 193
 - haemolytic uraemic syndrome, 193
 - pyelonephritis, 193
- reproductive choices, 181
- reproductive epidemiology trends, women healths and, 279
 - birth rates and fertility trends, 283
 - delivery complications, 290
 - delivery mode, 289
 - legal abortion trends, 281
 - maternal health aspects
 - alcohol, 292
 - blood pressure, 291
 - contraceptive pill usage, 292
 - ethnic populations health, 294
 - obesity, 291
 - postnatal depression and other psychological illness, 293
 - smoking, 292
 - maternities and estimated pregnancies, 279
 - maternities by
 - age, 284, 285
 - ethnic origin, 286, 287
 - marital status, 286
 - multiplicity, 288
 - National Statistics Socio-Economic group, 287, 288
 - parity, 286
 - reproductive health care information, 181
- research for preventing maternal deaths
 - caesarean section, 57
 - postpartum haemorrhage, 57
- respiratory system diseases, 195
 - acute respiratory distress syndrome, 190
 - asthma, 190
 - fibrosing alveolitis, 190
 - pneumonia, 190
- road traffic accidents during pregnancy
 - seat belts, requirement for, 208, 209
- Royal College of Obstetrics and Gynaecology (RCOG), 37
- Royal College of Pathologists (RCPATH), 224
- SADS. *See* sudden adult death syndrome
- Scottish Confidential Audit of Severe Maternal Morbidity
 - methods, 267
 - results, 268
- seat belts use in pregnancy
 - necessity of, 208, 209
 - recommendations for, 209
- severe maternal morbidity, 267, 270, 310
- sick pregnant women, recognizing for intensive care, 237
- social labelling of pregnant women, impact on cancer diagnosis, 202
- substance misuse
 - alcohol and drug, 162
 - causing maternal deaths, 50
 - dual therapy for, 179
 - effects on baby, 175
 - general practitioners, role of, 178
 - identification of, 178
 - midwifery practice issues, 259–260
 - substance misuse during the pregnancies, management of, 180
 - buprenorphine use, 181
 - methadone use, 181

- substandard care, 71, 306
 - causes of, 43
 - definitions of, 42
 - for amniotic fluid embolism, 100
 - for haemorrhage, 92
 - See *also* antenatal care
- sudden adult death syndrome (SADS), 147
 - bacterial endocarditis, 148
 - characteristics of deaths, 148
- sudden unexplained death in epilepsy (SUDEP), 187
- suicide
 - and infanticide, 169
 - cause of maternal death, 33, 156, 160–161
 - pathology report of, 231
 - profile (characteristics of pregnant women), 166
- suicide leading to maternal deaths
 - age at death, 161
 - education and training, 153
 - individual practitioners, 152
 - method of suicide (1997–2002), 160
 - service provision, 152
 - social characteristics, 161
- systolic blood pressure, 83

- Tenth International Classification of Diseases, Injuries and Causes of Death, The. See ICD10
- termination of pregnancy causing maternal deaths (examples), 106
- thromboembolism
 - as direct deaths cause, 243
 - causing maternal deaths, 61
 - 2000–02 report, 63
 - England and Wales 1952–84, 63
 - United Kingdom 1985–2002, 63
 - recommendations for preventing
 - RCOG guidelines on thromboprophylaxis, 61
 - venous thromboembolism (VTE) assessment, 61
 - risk factors for, 65
 - age, 66
 - anteartum deaths, 67
 - deaths after vaginal delivery, 67
 - delays in diagnosis, 68
 - family history, 66
 - immobilisation, 68
 - immobility, 66
 - management, 69
 - maternal weight, 66
 - obesity, 68
 - oral contraception, 67
 - previous history, 66
 - travel, 66
 - substandard care for, 71
 - See *also* amniotic fluid embolism
- thromboprophylaxis in obese women, 68
- thrombosis
 - 2000–02 report, 63
 - causing maternal deaths, 61
 - England and Wales 1952–84, 63
 - United Kingdom 1985–2002, 63
 - See *also* thromboembolism
- total fertility rate (TFR), 283
- toxicology
 - for epilepsy, 232
 - recommendations for, 232
- translation services problem, 47

- UK MMR
 - 1985–2002, 28
 - 2000–02, 29
 - See *also* maternal mortality rates
- uterine infection, risk of, 112. See *also* genital tract sepsis
- uterine rupture, maternal mortality from, 119

- vaginal delivery
 - deaths after, 67
 - genital tract sepsis after, 113
- ventricular-septal defect (VSD), 230
- viral infection
 - acute liver failure, 189
 - HIV/AIDS, 189
 - lactic acidosis, 189
- VTE (venous thromboembolism), 61. See *also* thromboembolism